

Title	ON CERTAIN ESTIMATES OF SINGULAR INTEGRALS USEFUL FOR EXTRAPOLATION (Harmonic Analysis and Nonlinear Partial Differential Equations)
Author(s)	SATO, SHUICHI
Citation	数理解析研究所講究録別冊 = RIMS Kokyuroku Bessatsu (2009), B14: 155-162
Issue Date	2009-11
URL	http://hdl.handle.net/2433/176890
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

ON CERTAIN ESTIMATES OF SINGULAR INTEGRALS USEFUL FOR EXTRAPOLATION

SHUICHI SATO

ABSTRACT. We consider several classes of singular integrals with rough kernels. We prove certain L^p estimates ($1 < p < \infty$) for the singular integrals. As an application, we can prove L^p boundedness of the singular integrals under a minimum size condition on their kernels via an extrapolation argument.

1. INTRODUCTION

Let a function Ω in $L^1(S^{n-1})$ satisfy

$$(1.1) \quad \int_{S^{n-1}} \Omega(\theta) d\sigma(\theta) = 0,$$

where S^{n-1} is the unit sphere in \mathbb{R}^n and $d\sigma$ is the Lebesgue surface measure on S^{n-1} . We assume $n \geq 2$. We consider singular integrals of the form:

$$(1.2) \quad \begin{aligned} T(f)(x) &= \text{p.v.} \int_{\mathbb{R}^n} f(x-y)K(y) dy, \\ K(x) &= h(|x|) \frac{\Omega(x')}{|x|^n}, \quad x' = x/|x|, \end{aligned}$$

where h is a function on $\mathbb{R}_+ = \{t \in \mathbb{R} : t > 0\}$.

Now we introduce two function spaces on S^{n-1} .

- (1) Let $L \log L(S^{n-1})$ denote the Zygmund class of all functions F on S^{n-1} satisfying

$$\int_{S^{n-1}} |F(\theta)| \log(2 + |F(\theta)|) d\sigma(\theta) < \infty.$$

- (2) The Hardy space $H^1(S^{n-1})$ is the space of functions $F \in L^1(S^{n-1})$ such that

$$\|F\|_{H^1(S^{n-1})} := \|P^+ F\|_{L^1(S^{n-1})} < \infty,$$

where $P^+ F$ is the radial maximal function defined as

$$P^+ F(\theta) = \sup_{0 \leq r < 1} \left| \int_{S^{n-1}} F(\omega) P_{r\theta}(\omega) d\sigma(\omega) \right|.$$

Here $P_{r\omega}(\theta)$ ($0 \leq r < 1$, $\omega, \theta \in S^{n-1}$) denotes the Poisson kernel:

$$P_{r\omega}(\theta) = c_n \frac{1 - r^2}{|r\omega - \theta|^n}.$$

It is known that $L \log L(S^{n-1})$ is a proper subspace of $H^1(S^{n-1})$.

We first assume that h is identically 1; so K is a homogeneous kernel. We write $T = T_\Omega$. Then,

$$(T_\Omega f)^\wedge(\xi) = m(\xi') \hat{f}(\xi),$$

where

$$m(\xi') = - \int_{S^{n-1}} \Omega(\theta) F(\xi', \theta) d\sigma(\theta),$$

$$F(\xi', \theta) = \left[i \frac{\pi}{2} \operatorname{sgn}(\langle \xi', \theta \rangle) + \log |\langle \xi', \theta \rangle| \right].$$

This implies, by Young's inequality, that $T_\Omega : L^2 \rightarrow L^2$ if $\Omega \in L \log L(S^{n-1})$. Also, by the method of rotations Calderón-Zygmund [4] proved that if Ω belongs to $L^1(S^{n-1})$ and is odd, then $T_\Omega : L^p \rightarrow L^p$ for all $1 < p < \infty$ and that if $\Omega \in L \log L(S^{n-1})$, then $T_\Omega : L^p \rightarrow L^p$ for all $1 < p < \infty$.

Furthermore, Coifman-Weiss [5], Connert [6] and Ricci-Weiss [15] proved that if $\Omega \in H^1(S^{n-1})$, then $T_\Omega : L^p \rightarrow L^p$ for all $1 < p < \infty$ by applying developed versions of the Calderón-Zygmund method of rotations. This is an improvement over the result of Calderón-Zygmund above, since $L \log L(S^{n-1})$ is a proper subspace of $H^1(S^{n-1})$.

Next, we see the case where h is not assumed to be a constant function. For $s \in [1, \infty)$, the space Δ_s is defined as

$$\Delta_s = \{h \text{ on } \mathbb{R}_+ : \|h\|_{\Delta_s} < \infty\},$$

where

$$\|h\|_{\Delta_s} = \sup_{j \in \mathbb{Z}} \left(\int_{2^j}^{2^{j+1}} |h(t)|^s dt/t \right)^{1/s}.$$

Here \mathbb{Z} denotes the set of all integers. Also, let $\Delta_\infty = L^\infty(\mathbb{R}_+)$. Then, we can easily see that $\Delta_s \subset \Delta_t$ if $s > t$.

If h is not constant, the method of rotations of Calderon-Zygmund does not work in general (see [17]), nevertheless, we have the following:

(a) (R. Fefferman [11]) If $h \in L^\infty$ and Ω satisfies a Lipschitz condition on S^{n-1} , then $T : L^p \rightarrow L^p$ for all $1 < p < \infty$.

J. Namazi [13] improved this result by replacing the condition on Ω with the L^q -condition:

(b) Suppose that $h \in L^\infty$ and $\Omega \in L^q(S^{n-1})$ for some $q > 1$. Then $T : L^p \rightarrow L^p$ for all $1 < p < \infty$.

J. Duoandikoetxea and J. L. Rubio de Francia [7] further improved this by replacing the condition on h with the Δ_2 condition:

(c) If $h \in \Delta_2$ and $\Omega \in L^q(S^{n-1})$ for some $q > 1$, then $T : L^p \rightarrow L^p$ for all $1 < p < \infty$. The L^q condition on Ω in (c) was relaxed by Fan and Pan [10] as follows:

(d) Suppose that $\Omega \in H^1(S^{n-1})$ and $h \in \Delta_s$ for some $s > 1$. Then $T : L^p \rightarrow L^p$ if $|1/p - 1/2| < \min(1/2, 1/s')$, $s' = s/(s-1)$.

We note the space $L^q(S^{n-1})$, $q > 1$, is a proper subspace of $H^1(S^{n-1})$ and when $s = 2$, the range of p in the conclusion of (d) is $(1, \infty)$.

Also, A. Al-Salman and Y. Pan [2] proved the following:

(e) If $\Omega \in L \log L(S^{n-1})$ and $h \in \Delta_s$ for some $s > 1$, then $T : L^p \rightarrow L^p$ for all

$1 < p < \infty$.

In (e) the condition on Ω is slightly stronger than that of (d), but the range of p shrinks to 2 as s approaches 1 in the conclusion of (d), while the range of p is always $(1, \infty)$ in the conclusion of (e), regardless the value of s .

Now, our first result is the following theorem.

Theorem 1. *Let T be as in (1.2) Suppose that $\Omega \in L^q(S^{n-1})$, $h \in \Delta_s$, $q, s \in (1, 2]$. Then*

$$\|T(f)\|_{L^p} \leq C_p(q-1)^{-1}(s-1)^{-1}\|\Omega\|_{L^q(S^{n-1})}\|h\|_{\Delta_s}\|f\|_{L^p}$$

for all $p \in (1, \infty)$, where the constant C_p is independent of q, s, Ω and h .

We are interested in this estimate when q and s are near 1. This estimate can be used to prove L^p boundedness of T by extrapolation of Yano under $L \log L$ condition for Ω and a certain condition for h . To state the condition for h , we need to introduce two more function spaces. For h on \mathbb{R}_+ and $a > 0$, let

$$L_a(h) = \sup_{j \in \mathbb{Z}} \int_{2^j}^{2^{j+1}} |h(r)| (\log(2 + |h(r)|))^a dr/r.$$

Define the class \mathcal{L}_a to be the space of all functions h satisfying $L_a(h) < \infty$. Then, we see that if $a < b$, $\mathcal{L}_b \subset \mathcal{L}_a$ and that

$$\bigcup_{s>1} \Delta_s \subsetneq \bigcap_{a>0} \mathcal{L}_a.$$

Also, for h on \mathbb{R}_+ and $a > 0$, let

$$N_a(h) = \sum_{m \geq 1} m^a 2^m d_m(h), \quad d_m(h) = \sup_{k \in \mathbb{Z}} 2^{-k} |E(k, m)|,$$

where

$$E(k, m) = \{r \in (2^k, 2^{k+1}] : 2^{m-1} < |h(r)| \leq 2^m\}$$

for $m \geq 2$ and

$$E(k, 1) = \{r \in (2^k, 2^{k+1}] : 0 < |h(r)| \leq 2\}.$$

We denote by \mathcal{N}_a the class of the functions h on \mathbb{R}_+ such that $N_a(h) < \infty$. Then, we can see that $N_a(h) < \infty$ implies $L_a(h) < \infty$. Conversely, if $L_{a+b}(h) < \infty$ for some $b > 1$, then $N_a(h) < \infty$.

Now, we can state an application of Theorem 1.

Theorem 2. *Let $\Omega \in L \log L(S^{n-1})$ and $h \in \mathcal{N}_1$. Then*

$$\|T(f)\|_{L^p} \leq C_p \|f\|_{L^p}$$

for all $p \in (1, \infty)$.

Theorem 2 follows from Theorem 1 and Yano's extrapolation (see [26]). Al-Salman-Pan [2] proved L^p boundedness of T under the condition that $\Omega \in L \log L(S^{n-1})$ and $h \in \Delta_s$ for some $s > 1$. Theorem 2 improves that result by replacing the assumption on h with the condition $h \in \mathcal{N}_1$. Here we recall that the condition $h \in \mathcal{N}_1$ follows if $h \in \mathcal{L}_a$ for some $a > 2$.

Proofs of Theorems 1 and 2 can be found in [18]. To prove Theorem 1 we apply the method of J. Duoandikoetxea and J. L. Rubio de Francia [7] involving the Littlewood-Paley theory. A new element of our proof is to apply a Littlewood-Paley decomposition adapted to a suitable lacunary sequence depending on q and s for which $\Omega \in L^q(S^{n-1})$ and $h \in \Delta_s$. The method of appropriately choosing a lacunary sequence has been already used in a different way from ours by A. Al-Salman and Y. Pan [2]. In the following sections, we shall see that Theorem 1 can extend to several classes of singular integrals.

2. SINGULAR RADON TRANSFORMS

Let

$$\begin{aligned} T(f)(x) &= \text{p.v.} \int_{\mathbb{R}^n} f(x - P(y))K(y) dy \\ &= \lim_{\epsilon \rightarrow 0} \int_{|y| > \epsilon} f(x - P(y))K(y) dy, \end{aligned}$$

where

$$K(y) = h(|y|)\Omega(y')|y|^{-n}, \quad y' = |y|^{-1}y,$$

$\Omega \in L^1(S^{n-1})$ satisfies (1.1), f is an appropriate function on \mathbb{R}^d (d may be different from n) and $P(y) = (P_1(y), P_2(y), \dots, P_d(y))$ is a polynomial mapping (each P_j is a real-valued polynomial on \mathbb{R}^n).

We assume that $P(-y) = -P(y)$. Then we have the following theorem.

Theorem 3. *Suppose that $\Omega \in L^q(S^{n-1})$, $h \in \Delta_s$, $q, s \in (1, 2]$. Then*

$$\|T(f)\|_{L^p(\mathbb{R}^d)} \leq C_p(q-1)^{-1}(s-1)^{-1}\|\Omega\|_{L^q(S^{n-1})}\|h\|_{\Delta_s}\|f\|_{L^p(\mathbb{R}^d)}$$

for all $p \in (1, \infty)$, where the constant C_p is independent of q, s, Ω and h . Also, the constant C_p is independent of polynomials P_j if we fix $\deg(P_j)$ ($j = 1, 2, \dots, d$).

By Theorem 3 and extrapolation we have the following result.

Theorem 4. *Let $\Omega \in L \log L(S^{n-1})$ and $h \in \mathcal{N}_1$. Then*

$$\|T(f)\|_{L^p(\mathbb{R}^d)} \leq C_p\|f\|_{L^p(\mathbb{R}^d)}$$

for all $p \in (1, \infty)$, where C_p is independent of polynomials P_j if the polynomials are of fixed degree.

Previous results are as follows.

- (a) (E. M. Stein [23]) If $h = 1$ and $\Omega \in C^1(S^{n-1})$, then $T : L^p \rightarrow L^p$ for all $1 < p < \infty$.
- (b) (D. Fan and Y. Pan [10]) Suppose that $\Omega \in H^1(S^{n-1})$ and $h \in \Delta_s$ for some $s > 1$. Then $T : L^p \rightarrow L^p$ if $|1/p - 1/2| < \min(1/2, 1/s')$.
- (c) (A. Al-Salman and Y. Pan [2]) Suppose that $\Omega \in L \log L(S^{n-1})$, $h \in \Delta_s$ for some $s > 1$ and $P(-y) = -P(y)$. Then $T : L^p \rightarrow L^p$ for all $1 < p < \infty$.

Theorem 4 improves the result (c) by replacing the assumption on h with $h \in \mathcal{N}_1$. For Theorems 3 and 4, see [18]. Relevant results can be found in [14], [24].

3. SINGULAR INTEGRALS ASSOCIATED WITH FUNCTIONS OF FINITE TYPE

We consider a singular Radon transform of the form:

$$T(f)(x) = \text{p.v.} \int_{B(0,1)} f(x - \Phi(y)) K(y) dy,$$

where $K(y) = \Omega(y')|y|^{-n}$, $\Omega \in L^1(S^{n-1})$, $y' = |y|^{-1}y$, $\Phi : B(0,1) \rightarrow \mathbb{R}^d$ is a smooth function, $B(0,1) = \{x \in \mathbb{R}^n : |x| < 1\}$. We assume that Ω satisfies (1.1) and that Φ is of finite type at the origin, where Φ is said to be of finite type at the origin if for any $\xi \in S^{d-1}$ there exists a multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$ such that $|\alpha| \geq 1$ and

$$\partial_x^\alpha \langle \Phi(x), \xi \rangle|_{x=0} \neq 0.$$

Then we have the following theorem.

Theorem 5. *Let $q \in (1, 2]$ and $\Omega \in L^q(S^{n-1})$. Then*

$$\|T(f)\|_{L^p(\mathbb{R}^d)} \leq C_p (q-1)^{-1} \|\Omega\|_{L^q(S^{n-1})} \|f\|_{L^p(\mathbb{R}^d)}$$

for all $p \in (1, \infty)$, where the constant C_p is independent of q and Ω .

By Theorem 5 and extrapolation, we can give a different proof for the following result of Al-Salman-Pan [2]:

Theorem A. *Suppose that $\Omega \in L \log L(S^{n-1})$. Then*

$$\|T(f)\|_{L^p(\mathbb{R}^d)} \leq C_p \|f\|_{L^p(\mathbb{R}^d)}$$

for all $p \in (1, \infty)$.

Relevant results are in [8]. See [20] for Theorem 5.

4. SINGULAR INTEGRALS ALONG SURFACES OF REVOLUTION

Let

$$\Gamma : [0, \infty) \rightarrow \mathbb{R}^m$$

be a continuous mapping satisfying $\Gamma(0) = 0$. We define a singular integral operator along the surface $(y, \Gamma(|y|))$ by

$$Tf(x, z) = \text{p.v.} \int_{\mathbb{R}^n} f(x - y, z - \Gamma(|y|)) K(y) dy$$

where $K(y) = h(|y|)\Omega(y')|y|^{-n}$. We assume that $\Omega \in L \log L(S^{n-1})$ satisfies (1.1).

Let

$$M_\Gamma g(z) = \sup_{R>0} R^{-1} \int_0^R |g(z - \Gamma(t))| dt.$$

We assume that $M_\Gamma : L^p(\mathbb{R}^m) \rightarrow L^p(\mathbb{R}^m)$ for all $p > 1$. An example of such Γ is a polynomial mapping. Γ may have infinite order contact with its tangent at the origin. Under this condition on Γ , we have the following theorem.

Theorem 6. *Suppose that $\Omega \in L^q(S^{n-1})$ for some $q \in (1, 2]$ and $h \in \Delta_s$ for some $s > 1$. Then,*

$$\|Tf\|_{L^p(\mathbb{R}^{n+m})} \leq C_p (q-1)^{-1} \|\Omega\|_q \|h\|_{\Delta_s} \|f\|_{L^p(\mathbb{R}^{n+m})}$$

if $|1/p - 1/2| < \min(1/s', 1/2)$, where the constant C_p is independent of q and Ω .

An extrapolation implies the following.

Theorem 7. Suppose $\Omega \in L \log L(S^{n-1})$ and $h \in \Delta_s$ for some $s > 1$. Then, T is bounded on $L^p(\mathbb{R}^{n+m})$ if $|1/p - 1/2| < \min(1/s', 1/2)$.

When $m = 1$ and Γ is a C^2 , convex, increasing function, Theorem 7 was proved by Al-Salman and Pan [2]. In that case, M_Γ is bounded on $L^p(\mathbb{R}^1)$ for all $p \in (1, \infty)$. See [19] for Theorems 6 and 7.

5. LITTLEWOOD-PALEY FUNCTIONS

We consider the Littlewood-Paley function on \mathbb{R}^n defined by

$$S_\psi(f)(x) = \left(\int_0^\infty |\psi_t * f(x)|^2 \frac{dt}{t} \right)^{1/2},$$

where ψ is in $L^1(\mathbb{R}^n)$, $\psi_t(x) = t^{-n}\psi(t^{-1}x)$. We assume that

$$\int_{\mathbb{R}^n} \psi(x) dx = 0.$$

One of the well-known sufficient conditions for L^p boundedness of S_ψ is the following:

Theorem B. Suppose that there exists $\epsilon > 0$ such that

$$|\psi(x)| \leq C(1 + |x|)^{-n-\epsilon},$$

$$\int_{\mathbb{R}^n} |\psi(x - y) - \psi(x)| dx \leq C|y|^\epsilon.$$

Then the operator S_ψ is bounded on $L^p(\mathbb{R}^n)$ for all $p \in (1, \infty)$.

This is due to Benedek, Calderón and Panzone [3]. It is known that the second assumption in Theorem B is not needed (see [16] and also [9]):

Theorem C. Suppose that

$$|\psi(x)| \leq C(1 + |x|)^{-n-\epsilon} \quad \text{for some } \epsilon > 0.$$

Then

$$S_\psi : L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n) \quad \text{for all } p \in (1, \infty).$$

Let

$$\psi(x) = |x|^{-n+1} \Omega(x') \chi_{(0,1]}(|x|), \quad x' = x/|x|,$$

where $\Omega \in L^1(S^{n-1})$ satisfies (1.1) and χ_E denotes the characteristic function of E . Define

$$\mu_\Omega(f) = S_\psi(f).$$

Then, $\mu_\Omega(f)$ is called the Marcinkiewicz integral (see Stein [22] and also Hörmander [12]). T. Walsh [25] proved the following result.

Theorem D. If $\Omega \in L(\log L)^{1/2}(S^{n-1})$, then

$$\mu_\Omega : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n).$$

Here $\Omega \in L(\log L)^{1/2}(S^{n-1})$ means

$$\int_{S^{n-1}} |\Omega(\theta)| [\log(2 + |\Omega(\theta)|)]^{1/2} d\sigma(\theta) < \infty.$$

Al-Salman, Al-Qassem, Cheng and Pan [1] extended Theorem D to all L^p ($1 < p < \infty$) spaces.

Theorem E. If $\Omega \in L(\log L)^{1/2}(S^{n-1})$, then

$$\mu_\Omega : L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n) \quad \text{for all } p \in (1, \infty).$$

We can give a different proof of Theorem E by extrapolation and the following result (see [21]).

Theorem 8. If $\Omega \in L^q(S^{n-1})$ for some $q \in (1, 2]$, we have

$$\|\mu_\Omega(f)\|_p \leq C_p (q-1)^{-1/2} \|\Omega\|_q \|f\|_p$$

for $p \in (1, \infty)$, where the constant C_p is independent of q and Ω .

REFERENCES

- [1] A. Al-Salman, H. Al-Qassem, L. C. Cheng and Y. Pan, *L^p bounds for the function of Marcinkiewicz*, Math. Res. Lett. **9** (2002), 697–700.
- [2] A. Al-Salman and Y. Pan, *Singular integrals with rough kernels in $L \log L(S^{n-1})$* , J. London Math. Soc. (2) **66** (2002), 153–174.
- [3] A. Benedek, A. P. Calderón and R. Panzone, *Convolution operators on Banach space valued functions*, Proc. Nat. Acad. Sci. U. S. A. **48** (1962), 356–365.
- [4] A. P. Calderón and A. Zygmund, *On singular integrals*, Amer. J. Math. **78** (1956), 289–309.
- [5] R. R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), 569–645.
- [6] W. Connnett, *Singular integrals near L^1* , Proc. Symp. Pure Math. **35**, Part 1 (1979), 163–165.
- [7] J. Duoandikoetxea and J. L. Rubio de Francia, *Maximal and singular integral operators via Fourier transform estimates*, Invent. Math. **84** (1986), 541–561.
- [8] D. Fan, K. Guo and Y. Pan, *Singular integrals along submanifolds of finite type*, Michigan Math. J. **44** (1997), 135–142.
- [9] D. Fan and S. Sato, *Remarks on Littlewood-Paley functions and singular integrals*, J. Math. Soc. Japan **54** (2002), 565–585.
- [10] D. Fan and Y. Pan, *Singular integral operators with rough kernels supported by subvarieties*, Amer. J. Math. **119** (1997), 799–839.
- [11] R. Fefferman, *A note on singular integrals*, Proc. Amer. Math. Soc. **74** (1979), 266–270.
- [12] L. Hörmander, *Estimates for translation invariant operators in L^p spaces*, Acta Math. **104** (1960), 93–139.
- [13] J. Namazi, *On a singular integral*, Proc. Amer. Math. Soc. **96** (1986), 421–424.
- [14] F. Ricci and E. M. Stein, *Harmonic analysis on nilpotent groups and singular integrals, I*, J. Func. Anal. **73** (1987), 179–194.
- [15] F. Ricci and G. Weiss, *A characterization of $H^1(\Sigma_{n-1})$* , Proc. Symp. Pure Math. **35**, Part 1 (1979), 289–294.
- [16] S. Sato, *Remarks on square functions in the Littlewood-Paley theory*, Bull. Austral. Math. Soc. **58** (1998), 199–211.
- [17] S. Sato, *Singular integrals and Littlewood-Paley functions*, Selected papers on differential equations and analysis, Translations. Series 2. **215** (2005), 57–78, American Mathematical Society, Providence, RI.
- [18] S. Sato, *Estimates for singular integrals and extrapolation*, Studia Math. **192** (2009), 219–233, arXiv:0704.1537v1 [math.CA].

- [19] S. Sato, *Estimates for singular integrals along surfaces of revolution*, to appear in J. Aust. Math. Soc.
- [20] S. Sato, *Estimates for singular integrals associated with manifolds of finite type and extrapolation*, preprint.
- [21] S. Sato, *Estimates for Littlewood-Paley functions and extrapolation*, Integr. equ. oper. theory **62** (2008), 429-440.
- [22] E. M. Stein, *On the functions of Littlewood-Paley, Lusin, and Marcinkiewicz*, Trans. Amer. Math. Soc. **88** (1958), 430-466.
- [23] E. M. Stein, *Problems in harmonic analysis related to curvature and oscillatory integrals*, Proceedings of International Congress of Mathematicians, Berkeley (1986), 196-221.
- [24] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton University Press, Princeton, NJ, 1993.
- [25] T. Walsh, *On the function of Marcinkiewicz*, Studia Math. **44** (1972), 203-217.
- [26] A. Zygmund, *Trigonometric Series*, 2nd ed., Cambridge Univ. Press, Cambridge, London, New York and Melbourne, 1977.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, KANAZAWA UNIVERSITY, KANAZAWA 920-1192, JAPAN

E-mail address: shuichi@kenroku.kanazawa-u.ac.jp